



# Major plant biome changes in the Primorye Region (Far East of Russia) during the Paleogene

Olesya V. Bondarenko<sup>1</sup>, Nadezhda I. Blokhina<sup>1\*</sup> & Torsten Utescher<sup>2,3</sup>

Olesya V. Bondarenko<sup>1</sup>  
e-mail: laricioxylon@gmail.com

Nadezhda I. Blokhina<sup>1\*</sup>  
e-mail: blokhina@biosoil.ru

Torsten Utescher<sup>2,3</sup>  
e-mail: t.utescher@uni-bonn.de

<sup>1</sup> Federal Scientific Center of the East Asia Terrestrial Biodiversity FEB RAS, Vladivostok, Russia

<sup>2</sup> Senckenberg Research Institute and Natural Museum, Frankfurt M., Germany

<sup>3</sup> Steinmann Institute, University of Bonn, Bonn, Germany

\* corresponding author

Manuscript received: 09.12.2018  
Review completed: 21.01.2019  
Accepted for publication: 29.01.2019  
Published online: 01.02.2019

Supplementary electronic information:  
[http://botsad.ru/media/aux/bp/BP\\_2019\\_8\\_1\\_bondarenko\\_e\\_suppl.xlsx](http://botsad.ru/media/aux/bp/BP_2019_8_1_bondarenko_e_suppl.xlsx)

## ABSTRACT

Major plant biome changes in Primorye (Far East of Russia) during the Paleogene are studied using the Integrated Plant Record vegetation analysis for the first time, based on palaeobotanical records. Palaeobotanical data employed for the reconstruction comprises 54 palaeofloras covering the early Paleocene to late Oligocene, a time-span of ca. 42 myr. Our data indicate the presence of more forested conditions in Primorye during the Paleogene than at present and demonstrate the major plant biome changes from the subtropical MMF/BLEF ecotone in the late Paleocene to the warm-temperate BLDF/MMF ecotone in the late Oligocene. The obtained major plant biome changes coincide in general with the climate cooling trend.

**Key words:** vegetation types, climate, spatial gradients, time series

## РЕЗЮМЕ

**Бондаренко О.В., Блохина Н.И., Утешер Т. Основные изменения растительных биомов Приморья (Дальний Восток России) в палеогене.** Основные изменения растительных биомов в Приморье (Дальний Восток России) в палеогене впервые изучены с использованием анализа растительности методом Integrated Plant Record, основанного на палеоботанических данных. Палеоботанические данные, примененные для реконструкции, включают 54 палеофлоры в интервале от раннего палеоцена до позднего олигоцена включительно, около 42 млн. лет. Наши данные указывают на наличие в Приморье более лесных условий в палеогене, чем в настоящее время, и демонстрируют основные изменения растительных биомов от субтропического экотона MMF/BLEF в позднем палеоцене до теплого умеренного экотона BLDF/MMF в позднем олигоцене. Полученные основные изменения растительных биомов в целом совпадают с тенденцией похолодания климата.

**Ключевые слова:** типы растительности, климат, пространственные градиенты, временные тренды

The Paleogene (65.5–23.03 Ma) globally was an interval of significant climatic and biotic reorganization (Akhmetiev 2004). It was a period of climate changes from greenhouse to icehouse conditions in the so-called “doubthouse” times marked by climatic cooling, rapid growth of the Antarctic ice sheet, and a supposed drop in atmospheric CO<sub>2</sub> levels leading to the Eocene-Oligocene Transition (EOT) event (Pagani et al. 2005, Dupont-Nivet et al. 2007, Eldrett et al. 2009, Pearson et al. 2009, Xiao et al. 2010, Abels et al. 2011).

Paleogene floras provide the potential for detailed interpretation of plant biomes and their response to global change such as warm climate thermal maxima and tectonic events. Therefore, the knowledge of the vegetation evolution during the Paleogene provides unique perspectives for the modeling of actual global changes and helps to probe into the integrated response of the Earth System to various driving forces (Collinson 1992, Zachos et al. 2008, Utescher et al. 2009).

The Primorye Region (or Primorye) is located in the south of the Russian Far East (RFE), on the coast of the Sea of Japan, bordering the Eurasian continent and Pacific Ocean. Located in the transitional zone “continent-ocean”, this territory constantly experiences the influence of circulation patterns related to the interaction of ocean, continental area, and atmosphere (Gerasimov 1969). Vegetation dynamics in the Paleogene can be reconstructed from the palaeobotanical record. Primorye is one of the Russian Far East regions, where large burials of Paleogene plants are concentrated, therefore the Paleogene flora of Primorye was a subject to extensive taxonomic studies. Results are presented in the works of Ablav (1974, 1977, 1978, 2000), Ablav & Akhmetiev (1977), Ablav et al. (2005, 2006), Akhmetiev (1973, 1974, 1988, 1993), Akhmetiev & Shevyreva (1989), Akhmetiev et al. (1973, 1978), Baskakova & Gromova (1979, 1984), Baskakova & Lepekhina (1990), M.D. Bolotnikova (1979), M.D. Bolotnikova & Sedykh (1987), T.N. Bolotnikova

(1988, 1989, 1993, 1994), Borsuk (1952), Gromova (1980), Klimova (1981, 1988), Klimova et al. (1977), Klimova & Tsar'ko (1989), Koshman (1964), Krassilov (1989a, b), Kundyshev & Petrenko (1987), Kundyshev & Verkhovskaya (1989), Mikhailov et al. (1989), Pavlyutkin (2007a, b, 2011), Pavlyutkin & Petrenko (1993, 1994, 1997, 2010), Pavlyutkin et al. (2006, 2012, 2014), Rybalko et al. (1980), Tashchi et al. (1996), Varnavskii et al. (1988), Verkhovskaya & Kundyshev (1989). However, these works are mainly devoted to the description of plant fossil remains, new fossil taxa, taxonomic diversity of Paleogene plants, or palaeofloristic aspects. Our knowledge of the evolution of the vegetation of Primorye in the Paleogene is still very poor and fragmentary, since most of the conclusions were made by different researchers on different materials (mainly based on few floras and/or limited territory) using different methods. These studies only provide some insights into local vegetation and its evolution, as well as indirectly about climate.


In the present detailed study we use the exceptionally rich palaeobotanical heritage of Primorye to reconstruct the Paleogene vegetation evolution of this region in space and time. Based on a total of 54 reasonably well-dated pollen and leaf floras we apply for the first time a consistent, timely method, the Integrated Plant Record vegetation analysis (IPR). The vegetation is reconstructed for seven stratigraphic levels and its dynamics is traced in a time-span of ca. 42 Myr, in total. Thus, our study not only unravels the regional vegetation evolution during the Paleogene but also promotes the understanding of the formation of modern vegetation in this region. Recently, Bondarenko et al (in print) provided a coherent palaeoclimate reconstruction for the Paleogene of Primorye based on the same palaeobotanical record. Thus, the present IPR-based biome reconstruction can be discussed in a palaeoclimatic context.

## MATERIAL AND METHODS

### Study area

The palaeobotanical record of Primorye studied herein originates from 19 basins (Fig. 1). The Paleogene strata of Primorye comprise volcanic and sedimentary deposits, unconformably overlying Mesozoic basement. The sedimentary facies includes fine- to coarse-grained continental clastics and intercalated lignites excavated in several active opencast mines. For some basins, mainly generated by extensional tectonics (Pavlovskii, Pushkinskii and Maksimovskii Basins), intercalated volcanoclastic layers and tholeiitic lava flows (Maksimovskii Basin: Takhobinskaya and Kuznetsovskaya formations) allow for radiometric dating of the strata (Table 1, Fig. 2). The sedimentary successions in the individual basins are characterized by numerous unconformities related to regional tectonics and phases of rifting and subsidence (Pavlyutkin & Petrenko 2010). When combining the strata of the individual basins a time-span of ca. 42 myr is represented, spanning the early Paleocene (Danian) to late Oligocene (Chattian).

The regional stratigraphic correlation chart for the basins (Fig. 2; adapted from Pavlyutkin & Petrenko 2010) is based on a variety of stratigraphic data obtained from radiometric dating, regional and inter-regional pollen zonation, as well



**Figure 1** Map showing the location of the Primorye Region within the Russian Far East and the location of the studied basins. Filled areas: the basins after Pavlyutkin & Petrenko (2010). 1 – Khasanskii, 2 – Ambinskii, 3 – Artemo-Tavrichanskii, 4 – Shkotovskii, 5 – Pushkinskii, 6 – Vanchinskii, 7 – Ivanovskii, 8 – Pavlovskii, 9 – Snegurovskii, 10 – Chernyshevskii, 11 – Zerkal'nenskii, 12 – Tur'erogskii, 13 – Krylovskii, 14 – Kemsii, 15 – Nizhnebikinskii, 16 – Amginskii, 17 – Maksimovskii, 18 – Svetlovodnenskii, 19 – Ozero Toni. The single floras are listed in Table 1, together with information on basin provenience, type of flora, stratigraphic age, method of dating, and references. The complete flora lists, assigned NLRs and their allocation to the components are given in Supplementary electronic information.

as lithological, palaeobotanical and vertebrate fauna correlations (Akhmetiev 1973, Varnavskii et al. 1988, Popov et al. 2005, Pavlyutkin & Petrenko 2010, Chatshin et al. 2013). The stratigraphic scheme has been tied to the International Stratigraphic Chart (Pavlyutkin & Petrenko 2010, Cohen et al. 2013) and allows for dating the flora-bearing horizons at the stage level (Fig. 2). For some of the floras stratigraphic ages are better constrained (cf. Table 1: radiometric datings for the Zanadvorovka, Gladkaya17, Kluch Stolbikova and Sobolevka floras).


### Floral record

The palaeobotanical record of Primorye is diverse and has been subject to extensive taxonomic studies (cf. Table 1

**Table 1.** Palaeofloras studied. P: palynofloras; L: leaf floras; IRC: inter-regional correlation based on paleobotany

Strat. level	Locality name (Lon/Lat)	Type of flora	No. of fossil taxa	Basin, formation	Age, method of dating	References
1. Late Oligocene	1. Zanadvorovka (131.4/43.2)	P	30	Ambinskii, analog of Pavlovskaya	Chatthian, K/Ar, (24.0±3.0 Ma), IRC	Pavlyutkin & Petrenko 2010, Popov et al. 2007
	2. Pushkinskii 9180 (131.5/43.3)	P L	30 28	Pushkinskii, Pavlovskaya	Chatthian, IRC	Pavlyutkin et al. 2012
	3. Pavlovka 9035-D (132.1/44.1)	P	35	Pavlovskii, Pavlovskaya	Chatthian, IRC	Pavlyutkin & Petrenko 2010
	4. Turii Rog 8 (131.6/45.1)	P	32	Tur'erogskii, analog of Pavlovskaya	Chatthian, IRC	Pavlyutkin & Petrenko 2010
	5. Luchegorsk 6212 (134.2/46.3)	P	32	Nizhnebikinskii, Upper Coal	Chatthian, IRC	Pavlyutkin & Petrenko 2010
2. Early Oligocene	6. Kraskino 9182/9196 (130.4/42.1)	P L	71 161	Khasanskii, Nizhnefatashinskaya	Rupelian, IRC	Pavlyutkin et al. 2014
	7. Pavlovka 9035-D (132.1/44.1)	P	37	Pavlovskii, Pavlovskaya	Rupelian, IRC	Pavlyutkin & Petrenko 2010
	8. Rettikhovka (132.4/44.1)	P L	36 48	Snegurovskii, analog of Pavlovskaya	Rupelian, IRC	Ablaev 1977, Akhmetiev et al. 1973, Klimova et al. 1977, Pavlyutkin & Petrenko 1994, 2010
	9. Voznovo 9206 (135.3/44.2)	P	79	Zerkal'nenskii, Voznovskaya	Rupelian, IRC	Pavlyutkin et al. 2014
	10. Tikhii Kluch (137.1/45.4)	L	49	Kemskii, Kizinskaya	Rupelian, IRC	Akhmetiev 1988, Rybalko et al. 1980, Varnavskii et al. 1988
	11. Amgu 9302 (137.4/45.6)	L	62	Amginskii, Grañatnenskaya	Rupelian, IRC	Akhmetiev 1988, Akhmetiev & Shyvareva 1989, Klimova 1981, 1988, Pavlyutkin et al. 2014
12. Maksimovka (137.5/46.1)	L	81	Maksimovskii, Maksimovskaya	Rupelian, IRC	Akhmetiev 1988, Rybalko et al. 1980, Varnavskii et al. 1988	
3. Late Eocene	13. Gladkaya 17 (130.5/42.4)	P L	41 33	Khasanskii, Khasanskaya	Priabonian, K/Ar (35.6–36.0 Ma), IRC	Chatshin et al. 2013, Gromova 1980, Pavlyutkin & Petrenko 1997, 2010, Pavlyutkin et al. 2006, 2014
	14. Tavrichanka 9142 (131.5/43.2)	P L	46 75	Artemo-Tavrichanskii, Ust'-davydovskaya	Priabonian, vertebrate fauna, IRC	Flerov et al. 1974, Pavlyutkin 2007a, Pavlyutkin & Petrenko 1993, 2010, Pavlyutkin et al. 2006, Yanovskaya 1954
	15. Shkotovo (132.2/43.2)	P	26	Shkotovskii, Upper Coal Fm.	Priabonian, IRC	Pavlyutkin & Petrenko 2010
	16. Ivanovka 610 (132.3/43.6)	P	50	Ivanovskii, analog of Pavlovskaya	Priabonian, IRC	Pavlyutkin & Petrenko 2010
	17. Pavlovka 9035-D (132.1/44.1)	P L	50 25	Pavlovskii, Pavlovskaya	Priabonian, IRC	Bolotnikova 1993, 1994, Pavlyutkin & Petrenko 2010, Varnavskii et al. 1988
	18. Svetlyi (135.1/44.1)	P L	14 22	Zerkal'nenskii, Svetlinskaya	Priabonian, IRC	Mikhailov et al. 1989, Pavlyutkin & Petrenko 2010
	19. Luchegorsk (134.2/46.3)	P L	61 56	Nizhnebikinskii, Bikinskaya	Priabonian, IRC	Ablaev et al. 2006, Bolotnikova & Sedykh 1987, Koshman 1964, Kundyshchev & Verkhovskaya 1989
	20. Salibeza (137.5/46.1)	L	48	Svetlovodnenskii, Salibežskaya	Priabonian, IRC	Klimova & Tsar'ko 1989, Varnavskii et al. 1988
4. Middle Eocene	21. Vol'no-Nadezhdinskoe (131.5/43.2)	P L	59 25	Artemo-Tavrichanskii, Nadezhdinskaya	Bartonian, IRC	Akhmetiev et al. 1978, Pavlyutkin 2007a, Pavlyutkin & Petrenko 1993, 2010
	22. Bolotnaya (131.1/43.2)	P L	45 75	Artemo-Tavrichanskii, Nadezhdinskaya	Lutetian, IRC	Ablaev 2000, Ablaev & Akhmetiev 1977, Akhmetiev 1973, Kundyshchev & Petrenko 1987, Pavlyutkin 2007, Pavlyutkin & Petrenko 2010
	23. Shkotovo (132.2/43.2)	P	31	Shkotovskii, Nadezhdinskaya	Middle Eocene, IRC	Baskakova & Gromova 1982
	24. Terekhovka (131.3/43.4)	P	44	Pushkinskii, Nadezhdinskaya	Middle Eocene, IRC	Pavlyutkin & Petrenko 2010
	25. Luchegorsk 540/541 (134.2/46.3)	P	33	Nizhnebikinskii, Luchegorskaya	Middle Eocene, IRC	Pavlyutkin & Petrenko 2010
5. Early Eocene	26. Tavrichanka 9142 (131.5/43.2)	P	58	Artemo-Tavrichanskii, Uglovskaya	Ypresian, IRC	Pavlyutkin and Petrenko 2010
	27. Smolyaninovo (132.3/43.2)	P L	56 42	Shkotovskii, Uglovskaya	Ypresian, IRC	Baskakova & Gromova 1979, 1984, Pavlyutkin & Petrenko 2010, Tashchi et al. 1996, Varnavskii et al. 1988, Verkhovskaya & Kundyshchev 1989
	28. Kluch Ugolnyi (134.1/43.3)	P L	19 20	Vanchinskii, analog of Uglovskaya	Ypresian, IRC	Chekryzhov et al. 2010, Pavlyutkin & Petrenko 2010
	29. Rettikhovka (132.4/44.1)	P	56	Snegurovskii, analog of Uglovskaya	Ypresian, IRC	Pavlyutkin & Petrenko 2010
	30. Arsen'evka (133.1/44.1)	P	57	Chernyshevskii, analog of Uglovskaya	Ypresian, IRC	Bolotnikova 1988
	31. Kluch Tuvanov (135.1/44.1)	L	62	Zerkal'nenskii, Tujanovskaya	Ypresian, IRC	Baskakova & Lepekhina 1990, Varnavskii et al. 1988
	32. Krylovskii 524 (133.4/45.1)	P	59	Krylovskii, Uglovskaya	Ypresian, IRC	Pavlyutkin & Petrenko 2010
	33. Luchegorsk 540/541 (134.2/46.3)	P L	58 12	Nizhnebikinskii, Lower Coal Fm.	Ypresian, IRC	Pavlyutkin & Petrenko 2010
34. Ozero Toni (138.3/47.4)	P L	32 44	Ozero Toni, Kizinskaya	Early Eocene, IRC	Oleinikov & Klimova 1977, Varnavskii et al. 1988	
6. Late Paleocene	35. Ustinovka (135.1/44.1)	L	25	Zerkal'nenskii, Tadushinskaya	Selandian, IRC	Pavlyutkin & Petrenko 2010
	36. Kluch Stolbikova (137.5/46.1)	L	14	Maksimovskii, Kuznetsovskaya	Thanetian, K/Ar (55.0 Ma), IRC	Pavlyutkin & Petrenko 2010, Varnavskii et al. 1988
	37. Kluch Kedrovyy (137.8/46.2)	P	37	Maksimovskii, Kedrovskaya	Thanetian, IRC	Varnavskii et al. 1988
7. Early Paleocene	38. Ustinovka (135.1/44.1)	L	31	Zerkal'nenskii, Bogopol'skaya	Danian, IRC	Ablaev et al. 2005, Krassilov 1989
	39. Sobolevka (137.5/46.1)	L	64	Maksimovskii, Takhobinskaya	Danian, K/Ar (64.0 Ma), IRC	Akhmetiev 1973, 1988, Borsuk 1952

**Note:** References and complete flora lists including Nearest Living Relatives used for vegetation analysis are given in Supplementary electronic information.



**Figure 2** Regional stratigraphic chart for the Paleogene sediments of some Cenozoic basins of Primorye considered in this study (modified from Pavlyutkin & Petrenko 2010), tied to the international standard (Cohen et al. 2013). Jagged boundary: unconformably overlying. Vertical stipple: gap in sedimentation. Details on the palaeofloras are given in Table 1. BgFm – Bogopol'skaya Formation, TkhFm – Takhobinskaya Formation (radiometric age 64.0 Ma by Akhmetiev 1973), TdFm – Tadushinskaya Formation, SuFm – Suvorovskaya Formation, KzFm – Kuznetsovskaya Formation (radiometric age 55.0 Ma by Varnavskii et al. 1988), UgFm – Uglovskaya Formation, TjFm – Tujanovskaya Formation, NdFm – Nadezhdinskaya Formation, KhFm – Khasanskaya Formation, UdFm – Ust'-davydovskaya Formation (radiometric age 35.6-36.0 Ma by Chatshin et al. 2013), SvFm – Svetlinskaya Formation, SbFm – Salibezskaya Formation, FtFm – Fatashinskaya Formation, PvFm – Pavlovskaya Formation, VzFm – Voznovskaya Formation, MsFm – Maksimovskaya Formation

for references). In this study, all palaeofloras considered here were carefully re-evaluated regarding the validity of taxonomic identifications and the Nearest Living Relatives (NLRs) of the fossil taxa, i.e. the taxonomical concept of the fossil materials and modern botanical affinity were carefully revised. In the present study, a total of 54 floras including 30 palynofloras (PF) and 24 leaf floras (LF) were studied with respect to palaeovegetation at seven stratigraphic levels. The floras cover a total time-span of ca. 42 myr, ranging from the early Paleocene (Danian) to late Oligocene (Chattian). The single floras are listed in Table 1, together with information on basin provenience, type of flora, stratigraphic age, method of dating, and references. Climate data for all floras are provided in Bondarenko et al. (in print).

### Integrated Plant Record vegetation analysis

The Integrated Plant Record vegetation analysis (IPR-vegetation analysis) is a semi-quantitative method first introduced by Kovar-Eder & Kvaček (2003) to assess zonal vegetation based on the fossil plant record (leaf, fruit, and

pollen assemblages). In order to employ the IPR, thirteen basic taxonomic-physiognomic groups, termed components, defined to reflect key ecological characteristics of an assemblage (Kovar-Eder & Kvaček 2003, 2007, Kovar-Eder et al. 2008, Teodoridis et al. 2011) are used: conifer component (CONIF), broad-leaved deciduous component (BLD), broad-leaved evergreen component (BLE), sclerophyllous component (SCL), legume-like component (LEG), zonal palm component (ZONPALM), arborescent fern component (ARBFERN), dry herbaceous component (D-HERB), mesophytic herbaceous component (M-HERB). Azonal components, i.e. azonal woody component (AZW), azonal non-woody component (AZNW) and aquatic component (AQUA). The component PROBLEMATIC TAXA includes elements with uncertain taxonomic-physiognomic affinity. For further analysis, all taxa (but not their abundances) of every single assemblage have to be assigned to those components and their relative proportions have to be calculated. The complete flora lists, assigned NLRs and their allocation to the components are given in Supplementary electronic information. The

number of taxa assigned to the components for each flora is given in Table 2.

To characterize zonal vegetation, the following proportions of components are regarded as relevant: (a) the proportion of the BLD, BLE, and SCL+LEG components of zonal woody angiosperms, where “zonal woody angiosperms” means sum of BLD+BLE+SCL+LEG+ZON-PALM+ARBFERN components; (b) the proportion of the ZONAL HERB (D-HERB+M-HERB) component of all zonal taxa, where “zonal taxa” means sum of the CONIF+BLD+BLE+SCL+LEG+ZONPALM+ARBFERN+D-HERB

+M-HERB components. The reliability of the results increases with increasing number of zonal taxa preserved. Ten zonal taxa are regarded as a minimum to perform this method (Kovar-Eder et al. 2008). Recently, Kovar-Eder & Teodoridis (2018) raised the former threshold to 15 zonal taxa for the application of the IPR-vegetation analysis. The proportions of the components were calculated for each flora and are given in Table 3.

Based on relative proportions of the components the following six zonal vegetation types are distinguished (Kovar-Eder & Kvaček 2007, Kovar-Eder et al. 2008): zonal

**Table 2.** Number of taxa allocated to the components (abbreviations are explained in text).

Stratigraphic level	Locality name	Type of flora	ZONAL										AZONAL			Problematic taxa
			CONIF	BLD	BLE	SCL	LEG	ZON PALM	ARB FERN	D-HERB	M-HERB	AZW	AZNW	AQU		
1. Late Oligocene	1. Zanadvorovka	P	5.5	15.14	2.81	1.28	0	0	0	0.2	0.4	4.44	0.2	0	0	
	2. Pushkinskii9180	P	5.5	15.14	2.81	1.28	0	0	0	0.2	0.4	4.44	0.2	0	0	
		L	0	17.16	0	0.66	2	0	0	0	0	6.16	0	2	0	
	3. Pavlovka9035-D	P	9.5	12.72	3.89	1.53	1	0	0	0.2	0.4	5.52	0.2	0	0	
	4. Turii Rog8	P	6.5	15.14	2.81	1.28	0	0	0	0.2	0.4	5.44	0.2	0	0	
5. Luchegorsk6212	P	8	14.69	2.86	1.53	0	0	0	0.2	0.2	4.49	0	0	0		
2. Early Oligocene	6. Kraskino9182/9196	P	12	25.62	6.29	1.78	1	0	0	0.78	4.48	12.67	1.28	2	0	
		L	20.5	74.63	28.63	1.5	10	0	0	0.5	2	20.63	1.5	1	0	
	7. Pavlovka9035-D	P	9.5	14.05	4.22	1.53	1	0	0	0.2	0.4	5.85	0.2	0	0	
	8. Rettikhovka	P	1.5	11.32	7.32	1.5	0	0.5	1	0.33	0.33	8.32	1.83	0	0	
		L	7	24.99	4.99	0.5	0	0	0	0	1.5	8.49	0.5	0	0	
	9. Voznovo9206	P	13.5	28.03	7.7	1.78	1	0	1	1.11	4.81	13.58	2.36	1	0	
		L	24.5	31.06	5.56	0.25	0	0	0	1	1	14.56	2	0	0	
	10. Tikhii Kluch	L	11	27.91	2.58	0.58	0	0	0	0.5	0.5	3.91	2	0	0	
	11. Amgu9302	L	16	25.14	5.81	0.33	1	0	0	0	1	11.64	0	1	0	
	12. Maksimovka	L	17	34.72	3.98	1.49	1	0	0	0.5	4.25	13.22	2.75	2	0	
	3. Late Eocene	13. Gladkaya17	P	6.5	12.73	6.07	1.41	0	0	0	0.33	0.33	9.23	1.33	0	0
			L	5	16.41	1.91	0.25	0	0	0	0	1	7.41	1	0	0
14. Tavrichanka9142		P	9.5	16.9	3.57	1.08	0	0	0	0.33	0.33	8.9	1.33	0	0	
		L	4	36.89	8.48	0.91	3	0	0	0.5	0.5	17.39	0.25	0	2	
15. Shkotovo		P	4	9.24	2.41	1.08	0	0	0	0.33	0.33	5.24	0.33	0	0	
16. Ivanovka610		P	12	14.07	7.24	1.58	0	0	0	0.33	0.83	9.07	0.83	0	0	
17. Pavlovka9035-D		P	11	16.51	4.01	0.5	1	0	1	0.99	2.69	8.51	2.19	1.5	0	
		L	0	16.66	2.83	0.83	0	0	0	0	0	3.66	0	1	0	
18. Svetlyi		P	0.5	4.83	4.33	1	0	0	0	0	0	3.33	0	0	0	
		L	5.5	3.33	0.33	0	0	0	0	0	5.5	4.83	2.5	0	0	
19. Luchegorsk		P	3.5	22.23	9.07	1.41	0	0	0	0.58	6.83	9.98	2.83	1.5	0	
		L	1	29.98	5.49	1.49	0	0	0	0	1	14.98	0	0	1	
20. Salibeza		L	9	13.32	2.32	0	0	0	0	0	7	10.32	2	4	0	
4. Middle Eocene	21. Vol'no-Nadezhdinskoe	P	8.5	20.81	6.98	1.83	0	0	0	0.33	2.33	9.81	2.33	0	0	
		L	1	11.66	2.08	0.58	0	0	0	0	2.25	6.16	0.75	0.5	0	
	22. Bolotnaya	P	10.5	16.65	3.82	1.83	0	0	0	0.33	1.33	9.15	0.33	0	0	
		L	0	46.05	9.47	0.83	1	0	0	0.25	1.75	14.3	0.25	0	1	
	23. Shkotovo	P	4.5	12.15	2.82	0.83	0	0	0	0	0	6.65	0	0	0	
	24. Terekhovka	P	8	17.47	7.64	1.28	0	0	0	0.2	0.4	7.77	0.2	0	0	
25. Luchegorsk540/541	P	7.5	11.85	4.52	1.33	0	0	0	0	0.2	5.35	0.2	0	0		
5. Early Eocene	26. Tavrichanka9142	P	9	19.73	7.4	1.58	0	0.5	1	0.33	1.83	10.23	1.33	0	0	
	27. Smolyaninovo	P	11.5	16.73	8.9	1.58	0	0	0	0.33	1.33	10.23	0.33	0	0	
		L	0	26.41	5.66	0	1	0	0	0	0.25	7.41	0.25	0	1	
	28. Kluch Ugolnyi	P	0.5	5.83	5.33	1	0	0	0	0	0	3.33	0	0	0	
		L	2	10	0.5	0	0	0	0	0	2	5.5	0	0	0	
	29. Rettikhovka	P	9.5	13.57	8.74	1.58	0	0	1	0.33	3.83	10.57	0.83	0	0	
	30. Arsen'evka	P	9	13.98	7.4	1.08	0	0	0	0.58	8.58	8.73	2.08	0.5	0	
	31. Kluch Tuyanov	L	11	24.57	3.82	0.5	0	0	0	0	4.25	15.57	2.25	0	0	
	32. Krylovskii524	P	10	18.63	8.3	1.78	0	0	1	1.03	3.73	11.43	1.03	0	0	
	33. Luchegorsk540/541	P	8.5	19.26	6.43	1.58	0	0	0	0.33	3.03	11.76	2.53	0.5	0	
		L	1	5.5	0	0	0	0	0	0	0	4.5	1	0	0	
	34. Ozero Toni	P	6	13.81	0.98	0.53	1	0	0	0.78	2.23	4.86	0.78	1	0	
		L	7	17.57	1.24	0.58	0	0	0	1	3	11.57	1	1	0	
	6. Late Paleocene	35. Ustinovka	L	1	14.33	0	0.33	0	0	0	0.33	2.83	5.33	0.83	0	0
36. Kluch Stolbikova		L	2	8.16	1.16	0	0	0	0	0	2.66	0	0	0	0	
37. Kluch Kedrovyyi		P	7.5	12.73	4.57	1.41	0	0	0	0.33	1.33	6.73	0.83	0.5	0	
7. Early Paleocene	38. Ustinovka	L	4.5	14.66	0.83	0.33	1	0	0	0	2	6.66	0	0	0	
	39. Sobolevka	L	10.5	26.65	5.07	0.58	0	0	0	0.5	5.75	13.15	0.75	0	0	

References and complete flora lists including Nearest Living Relatives used for vegetation analysis are given in Supplementary electronic information.

temperate to warm-temperate broad-leaved deciduous forests (broad-leaved deciduous forests, BLDF), zonal warm-temperate to subtropical mixed mesophytic forests (mixed mesophytic forests, MMF), zonal subtropical broad-leaved evergreen forests (broad-leaved evergreen forests, BLEF), zonal subtropical, subhumid sclerophyllous or microphyllous forests (subhumid sclerophyllous forests, ShSF), zonal xeric open woodlands (open woodland), and zonal xeric grasslands or steppe (xeric grassland). Teodoridis et al. (2011) additionally defined ecotones between the BLDF and MMF and the BLEF and MMF and recently, Kovar-Eder & Teodoridis (2018) defined ecotone between the MMF/ShSF.

## RESULTS

To apply IPR-vegetation analysis, 30 microfloras and 24 macrofloras were analyzed (Supplementary electronic information and Table 2). Based on the relative proportions of the components, three zonal vegetation types and two ecotones between them were revealed for the Paleogene of Primorye (Table 3).

The first (warmest) zonal vegetation type of subtropical broad-leaved evergreen forests (=broad-leaved evergreen forests, BLEF) was revealed from the late Eocene PF 18 and the early Eocene PF 28. These floras are characterized by low proportions of the BLD component (up to 48%), high proportions of the BLE component (42–44%), very low proportion of the SCL+LEG component (up to 10%), and absence of herbs.

The second zonal vegetation type of warm-temperate to subtropical mixed mesophytic forest (= mixed mesophytic forest, MMF) was revealed mainly based on PFs (from the late Paleocene to late Oligocene) and two LFs (the late Eocene and early Oligocene). The floras assigned to this zonal vegetation type are characterized by proportions of the BLD component from 62 to 75%, BLE component – 17–29%, SCL+LEG component – 5–13%, and 1–17% of herbs.


The third zonal vegetation type of temperate to warm-temperate broad-leaved deciduous forest (= broad-leaved deciduous forest, BLDF) was mainly reconstructed from LFs (from the early Paleocene to late Oligocene), and from the early Eocene PF 34. All these floras are characterized by high proportions of the BLD component (from 82 to 100%), low proportions of the BLE (0–16%), SCL+LEG (1–13%), and ZONAL HERB components (0–22%, except the late Eocene LF 18 constituting 37.5% of herbs).

The ecotones (MMF/BLEF and BLDF/MMF) are characterized by the transitional (intermediate) proportions of components in comparison with the corresponding zonal vegetation types (Table 3). The ecotone MMF/BLEF was revealed only for PFs from the early Eocene to early Oligocene and is characterized by 53–62% of BLD, 30–35% of BLE, 5–7% of SCL+LEG and 2–22% of herbs. The ecotone BLDF/MMF dominates in the late Oligocene and is characterized by 77–80% of BLD, 14–18% of BLE, 3–8% of SCL+LEG and 0–2% of herb components.

## DISCUSSION

### Comparison of micro- and macrofloras

The IPR claims to reconstruct vegetation independent from the type of flora, nevertheless, there is a distinct dif-



**Figure 3** Representation of IPR components in macro- and microfloras. Mean proportions calculated based on a total of 16 late Paleocene to late Oligocene localities yielding both, macro- and microrecords

ference in our results based on micro- and macrofloras. First of all, in the microfloras the number of zonal taxa is more stable in comparison with the macrofloras. The total number of all fossil taxa per site varies between 12 and 161 (Table 1). When ten zonal taxa are regarded as a minimum (Kovar-Eder et al. 2008), a mean number of zonal taxa of 34.8 for all the floras was sufficient to yield reliable results, except the LF 33 (6.5 zonal taxa only). When 15 zonal taxa are regarded as a minimum (Kovar-Eder & Teodoridis 2018) two microfloras and three macrofloras are insufficient to yield reliable results. The number of zonal taxa (Table 3) in each microflora ranged 10.66–58.93 (mean 32.3). The analysis of macrofloras was based on 6.5–137.76 (mean 37.8) zonal taxa.

The components used for calculating of the proportions also vary in the micro- and macrofloras (Table 2, Fig. 3). The CONIF component occurs in all microfloras ranging from 0.5 to 13.5, in the macrofloras – from 0 to 24.5 of zonal taxa. The BLD component is present in all micro- (4.8–28.6 zonal taxa) and macrofloras (3.3–74.6 zonal taxa). The BLE component in the microfloras varies from 1 to 9, whereas in the macrofloras – from 0 to 28.6 but mostly does not exceed 9.5 zonal taxa. The SCL component ranges from 0.5 to 1.8 in the microfloras and from 0 to 1.5 of zonal taxa in the macrofloras. The LEG component occurs rarely: 0–1 in microfloras, 0–2 zonal taxa in the macrofloras (except the early Oligocene LF 6 with 10 zonal taxa of the component). The ZONPALM component occurs only in some microfloras (0.5 zonal taxa) and is absent from the macrofloras. The ARBFERN component also occurs only in some microfloras (1 zonal taxa) and is absent from the macrofloras. The D-HERB component in the microfloras varies from 0.2 to 1.1, in the macrofloras – from 0 to 1 of zonal taxa. The M-HERB component is 0.2–8.6 zonal taxa in the microfloras and 0–7 zonal taxa in the macrofloras.

The ratio of each component in all micro- and macrofloras is given in Figure 3. As seen in the diagram (Fig. 3), IPR data are not independent from organ type regarded in each case. Leaf floras seem to be biased towards higher BLD and lower BLE proportions, microfloras have a lower proportion of LEG but a higher proportion of SCL components. As a result, the BLD forest biome is more frequently reconstructed based on leaves. This may have ta-

Table 3. Proportion of the components.

Stratigraphic level	Locality name	Palynofloras						Leaf floras						
		Number of zonal taxa	BLD prop. %	BLE prop. %	SCL+LEG prop. %	ZONAL HERB prop. %	Zonal vegetation type	Number of zonal taxa	BLD prop. %	BLE prop. %	SCL+LEG prop. %	ZONAL HERB prop. %	Zonal vegetation type	
1. Late Oligocene	1. Zanadvorovka	25.33	78.73	14.61	6.66	2.37	BLDF/MMF	—	—	—	—	—	—	
	2. Pushkinskiy180	25.33	78.73	14.61	6.66	2.37	BLDF/MMF	19.82	86.58	0.0	13.42	0.0	BLDF	
	3. Pavlovka9035-D	29.24	66.46	20.32	13.22	2.05	MMF	—	—	—	—	—	—	
	4. Turii Rog8	26.33	78.73	14.61	6.66	2.37	BLDF/MMF	—	—	—	—	—	—	
	5. Luchegorsk6212	27.48	76.99	14.99	8.02	1.45	BLDF/MMF	—	—	—	—	—	—	
2. Early Oligocene	6. Kraskino9182/9196	51.95	73.85	18.13	8.01	10.12	MMF	137.76	65.03	24.95	10.02	1.81	MMF	
	7. Pavlovka9035-D	30.9	67.55	20.29	12.16	1.94	MMF	—	—	—	—	—	—	
	8. Retikhovka	23.8	52.31	33.83	6.93	2.77	MMF/BLEF	38.98	81.99	16.37	1.64	3.85	BLDF	
	9. Voznovo206	58.93	70.94	19.49	7.04	10.04	MMF	63.37	84.24	15.08	0.68	3.16	BLDF	
	10. Tikhi Kluch	—	—	—	—	—	—	43.07	89.83	8.30	1.87	2.32	BLDF	
	11. Amgu9302	—	—	—	—	—	—	49.28	77.88	18.0	4.12	2.03	BLDF/MMF	
	12. Maksimovka	—	—	—	—	—	—	62.94	84.29	9.66	6.04	7.55	BLDF	
3. Late Eocene	13. Gladkaya17	27.37	62.99	30.03	6.98	2.41	MMF/BLEF	24.57	88.37	10.28	1.35	4.07	BLDF	
	14. Tavrichanka9142	31.71	78.42	16.57	5.01	2.08	BLDF/MMF	54.28	74.86	17.21	7.93	1.84	MMF	
	15. Shkotovo	17.39	72.58	18.93	8.48	3.79	MMF	—	—	—	—	—	—	
	16. Ivanovka610	36.05	61.47	31.63	6.90	3.22	MMF/BLEF	—	—	—	—	—	—	
	17. Pavlovka9035-D	37.7	71.72	17.42	6.52	9.76	MMF	20.32	81.99	13.93	4.08	0.0	BLDF	
	18. Svetvi	10.66	47.54	42.62	9.84	0.0	BLEF	14.66	90.98	9.02	0.0	37.52	BLDF	
	19. Luchegorsk	43.62	67.96	27.73	4.31	16.99	MMF	38.96	81.11	14.85	4.03	2.57	BLDF	
	20. Salibeza	—	—	—	—	—	—	31.64	85.17	14.83	0.0	22.12	BLDF	
	4. Middle Eocene	21. Vol'no-Nadezhdinskoe	40.78	70.26	23.56	6.18	6.52	MMF	17.57	81.42	14.52	4.05	12.81	BLDF
		22. Bolomaya	34.46	74.66	17.13	8.21	4.82	MMF	59.35	80.30	16.51	3.19	3.37	BLDF
23. Shkotovo		20.3	76.90	17.85	5.25	0.0	BLDF/MMF	—	—	—	—	—	—	
24. Terekhovka		34.99	66.20	28.95	4.85	1.71	MMF	—	—	—	—	—	—	
25. Luchegorsk540/541		25.4	66.95	25.54	7.51	0.79	MMF	—	—	—	—	—	—	
5. Early Eocene	26. Tavrichanka9142	41.37	65.31	24.49	5.23	5.22	MMF	—	—	—	—	—	—	
	27. Smolyanovo	40.37	61.48	32.71	5.81	4.11	MMF/BLEF	33.32	79.86	17.11	3.02	0.75	BLDF/MMF	
	28. Kluch Ugolnyi	12.66	47.94	43.83	8.22	0.0	BLEF	14.5	95.24	4.76	0.0	13.79	BLDF	
	29. Retikhovka	38.55	54.52	35.11	6.35	10.79	MMF/BLEF	—	—	—	—	—	—	
	30. Arsen'evka	40.62	62.24	32.95	4.81	22.55	MMF/BLEF	—	—	—	—	—	—	
	31. Kluch Tuyanov	—	—	—	—	—	—	44.14	85.05	13.22	1.73	9.63	BLDF	
	32. Krylovskiy524	44.47	62.71	27.94	5.99	10.70	MMF	—	—	—	—	—	—	
	33. Luchegorsk540/541	39.13	70.63	23.58	5.79	8.59	MMF	6.5	100.0	0.0	0.0	0.0	BLDF	
6. Late Paleocene	34. Ozero Toni	25.33	84.62	6.0	9.37	11.88	BLDF	30.39	90.61	6.39	2.99	13.16	BLDF	
	35. Ustinovka	—	—	—	—	—	—	18.82	97.75	0.0	2.25	16.79	BLDF	
7. Early Paleocene	36. Kluch Stolbikova	—	—	—	—	—	—	11.32	87.55	12.45	0.0	0.0	BLDF	
	37. Kluch Kedrovyy	27.87	68.04	24.42	7.54	5.96	MMF	—	—	—	—	—	—	
	38. Ustinovka	—	—	—	—	—	—	23.32	87.16	4.93	7.91	8.58	BLDF	
	39. Sobolevka	—	—	—	—	—	—	49.05	82.51	15.70	1.80	12.74	BLDF	

References and complete flora lists including Nearest Living Relatives used for vegetation analysis are given in Supplementary electronic information.

phonomic as well as edaphic reasons. Kovar-Eder & Kvaček (2007) assume that the differences of relative proportions of the components in the micro- versus the macrofloras are probably related to the lower taxonomic resolution of the pollen record and to the fact that pollen assemblages may be much more strongly influenced by long-distance transport than the leaf floras.

### Vegetation

As seen in the Table 3, the proportions calculated for determining of the zonal vegetation types differ in the micro- and macrofloras. The BLD proportions in the microfloras range from 52 to 84 %, in the macrofloras the proportions are higher (65–100 %). The BLE proportions in the microfloras are from 6 to 44 %, in the macrofloras – from 5 to 25 %. The SCL+LEG proportions are represented in all microfloras ranging from 5 to 13 %, while in the macrofloras the SCL+LEG proportions are 1–10 % and even in some macrofloras this component is absent. The ZONAL HERB proportions are represented in the microfloras up to 22 %, in the macrofloras – up to 37.5 %.

According to Kovar-Eder et al. (2008) and Teodoridis et al. (2011), with zonal herb proportions ranging from 0 to 22 %, except the late Eocene LF 18 constituting 37.5 % of herbs, no open woodland is reconstructed for the Primorye Region throughout the Paleogene based on the IPR-vegetation analysis. Moreover, according to Bondarenko et al. (2017), in the early Pleistocene the zonal herb proportion was ranging from 18 to 37 %. Thus, a close forest cover likely existed in Primorye throughout most of the Cenozoic. In Western Siberia, however, Popova et al. (2013) point out a steep declining trend of the arboreal component, already at the end of the Eocene.

Based on the microfloras, all three zonal vegetation types and two ecotones between them were revealed for the Paleogene of Primorye. The warmest zonal vegetation type (BLEF) was obtained only based on the microfloras and is represented by the early Eocene PF 28 and the late Eocene PF 18 (Zerkal'nenskii and Vanchinskii Basins correspondently, Fig. 4). The MMF zonal vegetation type dominated in Primorye during the Paleogene, as evident from the microfloras, and occurs from the late Paleocene to late Oligocene on the studied territory. The coldest zonal vegetation type (BLDF) was revealed only for the early Eocene PF 34 in the north of Primorye (Ozero Toni Basin, Fig. 4). The ecotone MMF/BLEF, reconstructed from microfloras, occurred from the early Eocene to early Oligocene, predominantly in the Eocene, in southern and middle part of Primorye, whereas the ecotone BLDF/MMF occurred from the middle Eocene to late Oligocene, but was predominant in the Oligocene.

The results obtained from the macrofloras are fundamentally different from the results based on microflora but may reflect more local conditions and wetland character of vegetation in the depocentres. The BLEF zonal vegetation type and the ecotone MMF/BLEF are not reconstructed from the macrofloras. The MMF zonal vegetation type was revealed only for the late Eocene LF 14 and early Oligocene LF 6 from the south of Primorye (Artemo-Tavrichanskii

and Khasanskii Basins correspondently). Unlike the microfloras, the BLDF zonal vegetation type is dominant in the macrofloras and occurred from the early Paleocene to late Oligocene on the studied territory. Moreover, only the ecotone BLDF/MMF was revealed for the early Eocene LF 27 and early Oligocene LF 11.

The modern vegetation of Primorye consists of mesophytic and xerophytic types (forest, wooded steppe, steppe, meadow, and swamp), which are in contact and form complex combinations (Kurentsova 1968). Our palaeobotanical data in general demonstrate a mesophytic forest (mixed broadleaved conifer forest with a diverse undergrowth) of subtropical to warm-temperate character in Primorye during the Paleogene while there is no evidence for xerophytic types and steppe.

The results obtained from the IPR-vegetation analysis of the Paleogene floral record of Primorye do not contradict the data obtained by other researchers. Ablaev et al. (2005), studying the late Paleocene LF 35, concluded that, in general, the territory under consideration was covered with coniferous-deciduous forests of warm-temperate type growing in the coastal region under a monsoon-type climate and this contributed to the development of a significant number of thermophilic elements here, compared to more continental areas.


Sedova (1957) and Flerov et al. (1974) infer for the south of Primorye in the early Eocene the presence of lush subtropical coastal vegetation, with the possible spread of mangroves there. However, Tashchi et al. (1996) question such conclusions and suggest warm temperate vegetation. According to present reconstruction, the early Eocene vegetation in Primorye was represented by a warm-temperate to subtropical vegetation according to the microfloras and mainly by a temperate to warm-temperate vegetation as reconstructed based on macrofloras (Table 4).

Pavlyutkin (2007) believes that the late Eocene LF 14 was probably formed in a warm-temperate climate with moderate humidity and represents a mesophytic, deciduous type, warm-temperate with some elements of more southern, subtropical origin. Our data also indicate a temperate to warm-temperate vegetation in the late Eocene according to the macrofloras, while the microfloras suggest a warm-temperate to subtropical vegetation (Table 4).

According to Pavlyutkin (2011) and Pavlyutkin et al. (2014), the early Oligocene LF 6 represents forest vegetation, mesothermal, transitional from warm-temperate to subtropical, mostly deciduous, although with significant participation of warmth-loving conifers and evergreen angiosperms. At the same time, the early Oligocene LF 9 and 11 represent deciduous, mesophilic, warm-temperate vegetation, with a minor participation of elements now found in mountain forests of the subtropics, some of which are evergreen (Pavlyutkin et al. 2014). This in general coincides with our reconstruction showing a warm-temperate to subtropical vegetation based on microfloras, and temperate to warm-temperate vegetation based on macrofloras, respectively, probably existing under specific local settings (Table 4).

There is evidence that in the Russian Far East, thermophilous vegetation persisted in the Neogene, even to the





**Figure 4** Proportions of the IPR components and corresponding zonal vegetation types in the Paleogene of Primorye based on microfossils: A – Early Oligocene, B – Late Oligocene, C – Middle Eocene, D – Late Eocene, E – Late Paleocene, F – Early Eocene. See explanations for abbreviations of IPR components and IPR biome types in the text

north of our study area. From early Neogene carpofloras of the eastern RFE, Popova et al. (2013) reconstructed a high diversity (40–50 %) of warm-temperate woody plants, including evergreen broad-leaved trees.

### Climate

It is known that changes in vegetation are associated with climate changes. Recently, the Paleogene climate dynamics

in Primorye are studied using the Coexistence Approach, based on the palaeobotanical records (Utescher et al. 2015, Bondarenko et al. in print). The climatic inferences obtained are consistent with independently-derived global trends, demonstrating general climate cooling throughout the Paleogene (Zachos et al. 2008). The data indicate that the Paleogene climate of Primorye was significantly warmer than present, in general, with the warmest conditions pre-

Table 4. Zonal vegetation types with climate data.

Stratigraphic level	Locality name	Palaenofloras				Leaf floras								
		Zonal vegetation type	MAT, C° <sup>1</sup>	MAP, mm <sup>1</sup>	Dominating zonal vegetation type	Mean MAT, C° <sup>2</sup>	Mean MAP, mm <sup>2</sup>	Zonal vegetation type	MAT, C° <sup>1</sup>	MAP, mm <sup>1</sup>	Dominating zonal vegetation type	Mean MAT, C° <sup>2</sup>	Mean MAP, mm <sup>2</sup>	
1. Late Oligocene	1. Zavadvorovka	BLDF/MMF	15	1188	BLDF/MMF	15.8	1213	—	—	—	BLDF	—	—	
	2. Pushkinskiĭ9180	BLDF/MMF	15	1188	—	—	—	—	14.4	1229	—	—	1229	
	3. Pavlovka9035-D	MMF	17.1	1278	—	—	—	—	—	—	—	—	—	
	4. Turĭ Rog8	BLDF/MMF	15	1132	—	—	—	—	—	—	—	—	—	
	5. Luchegorsk6212	BLDF/MMF	16.9	1278	—	—	—	—	—	—	—	—	—	
2. Early Oligocene	6. Krasino9182/9196	MMF	16.9	1163	MMF	16.8	1296	MMF	16.3	1118	BLDF	15.6	1150	
	7. Pavlovka9035-D	MMF	17.4	1278	—	—	—	—	—	—	—	—	—	
	8. Retikhovka	MMF/BLEF	16.9	1422	—	—	—	—	14.8	1329	BLDF	—	—	
	9. Voznovo9206	MMF	16	1323	—	—	—	—	15.4	1134	BLDF	—	—	
	10. Tikhiĭ Kluch	—	—	—	—	—	—	—	15.2	1059	BLDF/MMF	—	—	
	11. Amgu9302	—	—	—	—	—	—	—	15.4	1137	BLDF	—	—	
	12. Maksimovka	—	—	—	—	—	—	—	16.5	1126	BLDF	—	—	
3. Late Eocene	13. Gladkaya17	MMF/BLEF	16.9	1554	MMF	16.9	1432	MMF	17.6	1307	BLDF	17	1171	
	14. Tavrichanka9142	BLDF/MMF	16.9	1554	—	—	—	—	17.6	1227	—	—	—	
	15. Shkotovo	MMF	17.6	1278	—	—	—	—	—	—	—	—	—	
	16. Ivanovka610	MMF/BLEF	18.2	1404	—	—	—	—	—	—	—	—	—	
	17. Pavlovka9035-D	MMF	16.5	1373	—	—	—	—	17.1	1044	BLDF	—	—	
	18. Svetlyĭ	BLEF	16.3	1596	—	—	—	—	14.8	1158	BLDF	—	—	
	19. Luchegorsk	MMF	16.1	1265	—	—	—	—	17.2	1198	BLDF	—	—	
	20. Salibeza	—	—	—	—	—	—	—	17.5	1095	BLDF	—	—	
	4. Middle Eocene	21. Vol'no-Nadezhdinskoe	MMF	16.9	1554	MMF	17.2	1295	MMF	17.8	1172	BLDF	17.9	1228
		22. Bolotnaya	MMF	17.4	1170	—	—	—	—	18	1284	—	—	—
23. Shkotovo		BLDF/MMF	17.5	1070	—	—	—	—	—	—	—	—	—	
24. Terekhovka		MMF	17.3	1404	—	—	—	—	—	—	—	—	—	
25. Luchegorsk540/541		MMF	17.1	1278	—	—	—	—	—	—	—	—	—	
26. Tavrichanka9142		MMF	17.3	1554	MMF/BLEF	17.6	1338	—	—	—	BLDF	16.2	1162	
27. Smolyaninovo		MMF/BLEF	18.2	1295	—	—	—	—	17.5	1315	BLDF/MMF	—	—	
5. Early Eocene	28. Kluch Ugolnyi	BLEF	18.2	1422	—	—	—	—	17.2	1122	BLDF	—	—	
	29. Retikhovka	MMF/BLEF	18.2	1351	—	—	—	—	—	—	—	—	—	
	30. Arsen'evka	MMF/BLEF	18.2	1404	—	—	—	—	—	—	—	—	—	
	31. Kluch Tuyanov	—	—	—	—	—	—	—	15.9	1170	BLDF	—	—	
	32. Krylovskii524	MMF	16.6	1278	—	—	—	—	—	—	—	—	—	
	33. Luchegorsk540/541	MMF	16.6	1278	—	—	—	—	15.2	1159	BLDF	—	—	
	34. Ozero Torĭ	BLDF	17.4	1126	—	—	—	—	15.1	1047	BLDF	—	—	
6. Late Paleocene	35. Ustinovka	—	—	—	MMF	17.6	1158	—	16.9	1045	BLDF	17.1	1127	
	36. Kluch Stolbikova	—	—	—	—	—	—	—	17.3	1210	BLDF	—	—	
	37. Kluch Kedrovyyĭ	MMF	17.6	1158	—	—	—	—	—	—	—	—	—	
7. Early Paleocene	38. Ustinovka	—	—	—	—	—	—	—	18.4	1045	BLDF	16.4	1169	
	39. Sobolevka	—	—	—	—	—	—	—	14.5	1293	BLDF	—	—	

References and complete flora lists including Nearest Living Relatives used for vegetation analysis are given in Supplementary electronic information. <sup>1</sup> Means of Coexistence Intervals (Bondarenko et al., in print); <sup>2</sup> Means by substage calculated from <sup>1</sup>

vailing throughout the Eocene and in the southeast of the study area. Negligible Paleogene temperature gradients over Primorye are related to the global pattern revealing shallow temperature gradients in general (Greenwood & Wing 1995, Utescher et al. 2011), and specific regional aspects (Bondarenko et al. in print). The precipitation reconstruction points to conditions considerably wetter than at present (Utescher et al. 2014). A distinct increase in mean annual precipitation was reconstructed for the study area for the early Eocene and moderately declined throughout the Oligocene (Utescher et al. 2014). The regional rainfall pattern fundamentally differed from modern conditions, and this holds for all studied variables. The inland region and the south of Primorye were significantly more humid than today. The Paleogene pattern was possibly related to a monsoon-type circulation and enhanced landward flow of humid air masses, possibly due to an overall flatter morphology of the East Asian coastal areas (Bondarenko et al. in print) and a weak Siberian High that plays a critical role in the regional atmospheric circulation patterns (Popova et al. 2012).

According to Bondarenko et al. (in print), the mean MAT values based on the microfloras demonstrate the direct cooling trend (Table 4). In general, the obtained major plant biome changes (the dominating zonal vegetation type) in Primorye during the Paleogene, from the subtropical MMF/BLEF ecotone in the late Paleocene to the warm-temperate BLDF/MMF ecotone in the late Oligocene, as reconstructed based on the microfloras (Table 4) coincides with this cooling trend.

At the same time, the mean MAT values reconstructed from the macrofloras, having lower MAT values in general, show warm peaks in the late Paleocene and middle Eocene (Table 4) corresponding to the Paleocene–Eocene Thermal Maximum (PETM) and climatic optimum in the middle Eocene (MECO). However, despite the larger amplitude of temperatures and the presence of warm peaks, the macrofloras do not reflect major changes in vegetation, and show only one dominating type of temperate to warm-temperate forest – the BLDF (Table 4). Most likely, this can be explained by the fact that the leaves of deciduous plants are more likely to be buried than those of evergreen plants, while pollen show their presence more reliably (Krassilov 1972).

According to Akhmetiev (2004), Primorye belonged to a warm-temperate, humid climate zone during the Paleocene. In the early Eocene, this warm-temperate, humid zone was significantly reduced in size and covered mainly the polar region. At the transition from a warm-temperate climate to a subtropical zone in East Asia, especially along the Pacific Coast, a wide ecozone with deciduous and evergreen elements established. Deciduous elements in the conditions of the dissected relief of the oceanic margin penetrated far to the south, and in the zones of coastal evergreen plants, including palm trees, to the north (Akhmetiev 2004). In the middle Eocene, according to Akhmetiev (2004), the zone of temperate and warm-temperate humid climate, which existed from the beginning of the Paleogene, comprised the circumpolar region of the Northern Hemisphere, although its southern boundary shifted somewhat to the south, compared with its position in the Early Eocene. At the beginning of the middle Eocene (Lutetian)

in eastern Eurasia, a zone of mesophilic subtropical floras (a zone of humid and variable-humid (seasonal) subtropical climate) encompassed the very south of Primorye.

Paleogene temperature records are available for northeast China and thus from the closer neighborhood of our study area (Quan et al. 2012b). Features in common are the declining trend from the Paleocene to the early Eocene (CMMT), and a very warm late Eocene with subsequent distinct cooling (MAT, CMMT). The very warm late Eocene is not reflected in the global climate evolution and thus may represent a regional signal, possibly related to the initial opening of the Sea of Japan. According to Quan et al. (2012b), among the estimated terrestrial temperature parameters reconstructed from a sequence exposed in the Fushun open cast, MAT slightly changed with an overall declining trend in the Eocene. However, the winter temperature dramatically decreased from the middle Eocene to the late Eocene, while the summer temperature remained almost the same throughout the Eocene. All these data are within the range of the palaeoclimate reconstruction by Bondarenko et al. (in print) or slightly higher (CMMT), respectively, which coincides with the latitude of Fushun, lower by several degrees.

Quan et al. (2012a) demonstrate that in the Eocene, monsoonal climate probably had already been established over China, judging from the presence of apparent seasonality of both temperature and precipitation. In addition, Quan et al. (2012a) point out that, with a slight declining trend of MAT during the Eocene, the winter temperature substantially dropped in tropical southern China during the middle to late Eocene interval. According to Quan et al. (2012a), this might have been related to the development of a weak Eocene Kuroshio Current in the southwestern Pacific, and/or a significantly enhanced palaeo-winter monsoon from Siberia.

## CONCLUSIONS

The exceptionally rich palaeobotanical record of Primorye holds the key for reconstructing the detailed Paleogene regional vegetation evolution in space and time. The high diversity of the palaeofloras and up-to-date taxonomy result in useful vegetation interpretations. Our vegetation maps for the first time allow quantifying major plant biome changes in space and time on the Pacific side of Eurasia over the past 42 myr. Our palaeobotanical data indicate more forested conditions within the studied territory during the Paleogene than at present and in general prove the existence of a mesophytic forest (mixed broadleaved conifer forest with a diverse undergrowth) of subtropical to warm-temperate character in Primorye through the Paleogene. Moreover, the obtained major plant biome changes in Primorye from the subtropical MMF/BLEF ecotone in the late Paleocene to the warm-temperate BLDF/MMF ecotone in the late Oligocene, as reconstructed based on the microfloras, coincide with the coeval, globally observed cooling trend.

## ACKNOWLEDGEMENTS

The study was supported by the Russian Foundation for Basic Research (project no. 16-04-01241) to Nadezhda I. Blokhina. This work is a contribution to NECLIME network (Neogene Climate Evolution in Eurasia).

## LITERATURE CITED

- Abels, H.A., G. Dupont-Nivet, G. Xiao, R. Bosboom & W. Krijgsman 2011. Step-wise change of Asian interior climate preceding the Eocene – Oligocene Transition (EOT). *Paleogeography, Paleoclimatology, Paleoecology* 299:399–412.
- Ablaev, A.G. 1974. Additional data on the morphology of fossil leaves of *Liquidambar* of the Far East. In: *Issues of biostratigraphy of the Soviet Far East* (E.A. Kireeva & A.G. Ablaev, eds), pp. 113–117, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Аблаев А.Г. 1974. Дополнительные данные по морфологии ископаемых листьев ликвидамбра Дальнего Востока // Вопросы биостратиграфии советского Дальнего Востока / под ред. Е.А. Киреевой, А.Г. Аблаева. Владивосток: ДВНЦ АН СССР. С. 113–117]
- Ablaev, A.G. 1977. Flora of coal-bearing Neogene deposits of Primorye. In: *Stratigraphy of the Cenozoic deposits of the Far East* (V.G. Varnavskii, ed.), pp. 54–58, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Аблаев А.Г. 1977. Флора угленосных неогеновых отложений Приморья // Стратиграфия кайнозойских отложений Дальнего Востока / под ред. В.Г. Варнавского. Владивосток: ДВНЦ АН СССР. С. 54–58].
- Ablaev, A.G. 1978. *Geology and history of the coastal regions of the Sea of Japan*. Nauka, Moscow, 192 pp. (in Russian). [Аблаев А.Г. 1978. Геология и история побережья Японского моря. Москва: Наука. 192 с.].
- Ablaev, A.G. 2000. *Paleogene biostratigraphy of the coastal region in south Primor'e*. Dal'nauka, Vladivostok, 117 pp. (in Russian). [Аблаев А.Г. 2000. Биостратиграфия палеогена побережья юга Приморья. Владивосток: Дальнаука. 117 с.].
- Ablaev, A.G. & M.A. Akhmetiev 1977. Bolotninskaya Miocene flora of the Southern Primorye and the role of thermophilic elements in its composition. *Paleontologicheskii Zhurnal* 1:134–141 (in Russian). [Аблаев А.Г., Ахметьев М.А. 1977. Болотнинская миоценовая флора Южного Приморья и роль в её составе термофильных элементов // Палеонтологический журнал. № 1. С. 134–141].
- Ablaev, A.G., C.S. Li, I.V. Vassiliev & Y.F. Wang 2005. *Paleogene of the eastern Sikhote-Alin*. Dal'nauka, Vladivostok, 95 pp. (in Russian). [Аблаев А.Г., Ли Чен-Сен, Васильев И.В., Ван Ю-Фей 2005. Палеоген Восточного Сихотэ-Алиня. Владивосток: Дальнаука. 95 с.].
- Ablaev, A.G., C.S. Li & Y.F. Wang 2006. *Paleogene of the Bikin-Ussuri Basin of sedimentation (Lower-Bikin Depression)*. Dal'nauka, Vladivostok, 75 pp. (in Russian). [Аблаев А.Г., Ли Чен-Сен, Ван Ю-Фей 2006. Палеоген Бикино-Уссурийского бассейна седиментации (Нижне-бикинская впадина). Владивосток: Дальнаука. 75 с.].
- Akhmetiev, M.A. 1973. Paleocene and Eocene flora of the southern Far East of the USSR and adjacent countries and their stratigraphic position. *Sovetskaya geologiya* 7:14–29 (in Russian). [Ахметьев М.А. 1973. Палеоценовые и эоценовые флоры юга Дальнего Востока СССР и сопредельных стран и их стратиграфическое положение // Советская геология. № 7. С. 14–29].
- Akhmetiev, M.A. 1974. Oligocene and Miocene flora of the south of the Far East of the USSR as an indicator of the climatic situation. *Izvestiya Akademii nauk SSSR, Ser. Geologicheskaya* 4: 134–143 (in Russian). [Ахметьев М.А. 1974. Олигоценовые и миоценовые флоры юга Дальнего Востока СССР как показатель климатической обстановки // Известия АН СССР, Сер. Геологическая. № 4. С. 134–143].
- Akhmetiev, M.A. 1988. *Cenozoic flora of eastern Sikhote-Alin*. Geologicheskii institut Akademii nauk SSSR, Moscow, 48 pp. (in Russian). [Ахметьев М.А. 1988. Кайнозойские флоры восточного Сихотэ-Алиня. Москва: ГИН АН СССР. 48 с.].
- Akhmetiev, M.A. 1993. *Phytostratigraphy of Paleogene and Miocene continental deposits of Boreal Asia* (Trudy Geologicheskogo instituta Rossiiskoi akademii nauk, Vol. 475). Nauka, Moscow, 143 pp. (in Russian). [Ахметьев М.А. 1993. Фитостратиграфия континентальных отложений палеогена и миоцена внетропической Азии. (Труды ГИН РАН. Вып. 475). Москва: Наука. 143 с.].
- Akhmetiev, M.A. 2004. The Paleocene and Eocene global climate. Paleobotanical evidences. In: *Climate in the Epochs of Major Biospheric Transformations* (Trudy Geologicheskogo instituta RAN, Vol. 550) (M.A. Semikhatov & N.M. Chumakov, eds), pp. 10–43, Geologicheskii institut Rossiiskoi akademii nauk, Moscow (in Russian). [Ахметьев М.А. 2004. Климат земного шара в палеоцене и эоцене по данным палеоботаники // Климат в эпохи крупных биосферных перестроек (Труды ГИН РАН. Вып. 550) / под ред. М.А. Семихатова, Н.М. Чумакова. Москва: ГИН РАН. С. 10–43].
- Akhmetiev, M.A., M.D. Bolotnikova, G.M. Bratseva & V.A. Krassilov 1978. Stratigraphy and paleofloristics of the reference section of the Cenozoic of the Southern Primorye. *Izvestiya Akademii nauk SSSR, Ser. Geologicheskaya* 4:61–75 (in Russian). [Ахметьев М.А., Болотникова М.Д., Братцева Г.М., Красилов В.А. 1978. Стратиграфия и палеофлористика опорного разреза кайнозой Южного Приморья // Известия АН СССР, Сер. Геологическая. № 4. С. 61–75].
- Akhmetiev, M.A., G.M. Bratseva & R.S. Klimova 1973. On the age analogues of the Korean layers with Engelhardia in Primorye. *Doklady Akademii nauk SSSR* 209(1):167–170 (in Russian). [Ахметьев М.А., Братцева Г.М., Климова Р.С. 1973. О возрастных аналогах энгельгардиевых слоев Корен в Приморье // Доклады АН СССР. Т. 209. № 1. С. 167–170].
- Akhmetiev, M.A. & N.A. Shyvareva 1989. Fossil gymnosperms of the Amgu (East Sikhote-Alin). In: *Paleofloristics and stratigraphy of the Phanerozoic* (M.A. Akhmetiev, ed.), pp. 104–117, Geologicheskii institut Akademii nauk SSSR, Moscow (in Russian). [Ахметьев М.А., Шыварёва Н.А. 1989. Ископаемые голосеменные Амгу (Восточный Сихотэ-Алинь) // Палеофлористика и стратиграфия фанерозоя / под ред. М.А. Ахметьева. Москва: ГИН АН СССР. С. 104–117].
- Baskakova, L.A. & N.S. Gromova 1979. Biostratigraphic dismemberment of the Uglvskii horizon according to palynological data. In: *Paleontology and stratigraphy of the Far East* (B.V. Poyarkov, ed.), pp. 109–114, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Баскакова Л.А., Громова Н.С. 1979. Биостратиграфическое расчленение угловского горизонта по палинологическим данным // Палеонтология и стратиграфия Дальнего Востока / под ред. Б.В. Пояркова. Владивосток: ДВНЦ АН СССР. С. 109–114].
- Baskakova, L.A. & N.S. Gromova 1982. Phytostratigraphic dismemberment of Paleogene sediments of the South-Western Primorye. *Sovetskaya geologiya* 11: 68–78. (in Russian). [Баскакова Л.А., Громова Н.С. 1982. Фитостратиграфическое расчленение палеогеновых отложений Юго-Западного Приморья // Советская геология. № 11. С. 68–78].
- Baskakova, L.A. & N.S. Gromova 1984. Stratigraphy of Smolyaninovskii coal mine in the Southern Primorye.

- In: *Materials on stratigraphy and paleogeography of East Asia* (A.G. Ablaev, ed.), pp. 59–69, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Баскакова Л.А., Громова Н.С. 1984. Стратиграфия Смоляниновского угольного разреза в Южном Приморье // Материалы по стратиграфии и палеогеографии Восточной Азии / под ред. А.Г. Аблаева. Владивосток: ДВНЦ АН СССР. С. 59–69].
- Baskakova, L.A. & V.G. Lepekina. 1990. New data on the phytostратigraphy of the Paleogene of the Zerkal'enskii Basin. In: *New data on stratigraphy of the Far East and the Pacific Ocean* (A.G. Ablaev, ed.), pp. 52–60, Dal'nevostochnoe otdelenie Rossiiskoi akademii nauk, Vladivostok (in Russian). [Баскакова Л.А., Лепёхина В.Г. 1990. Новые данные по фитостратиграфии палеогена Зеркаленской впадины // Новые данные по стратиграфии Дальнего Востока и Тихого океана / под ред. А.Г. Аблаева. Владивосток: ДВО РАН. С. 52–60].
- Bolotnikova, M.D. 1979. *Spore-pollen complexes of Tertiary sediments of the western coast of the Sea of Japan*. Nauka, Moskva, 196 pp. (in Russian). [Болотникова М.Д. 1979. Спорово-пыльцевые комплексы третичных отложений западного побережья Японского моря. Москва: Наука. 196 с.].
- Bolotnikova, M.D. & A.K. Sedykh 1987. New data on the stratigraphy of coal-bearing sediments of the Nizhnebikinskii Basin. In: *Palinology of the East of the USSR* (V.S. Markevich, ed.), pp. 41–52, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Болотникова М.Д., Седых А.К. 1987. Новые данные по стратиграфии угленосных отложений Нижнебикинской впадины // Палинология Востока СССР / под ред. В.С. Маркевича. Владивосток: ДВНЦ АН СССР. С. 41–52].
- Bolotnikova, T.N. 1988. Palynological characteristics and age of coal-bearing deposits of the Chernyshevskii brown coal field (Southern Primorye). *Tikhoookeanskaya geologiya* 4: 101–105 (in Russian). [Болотникова Т.Н. 1988. Палинологическая характеристика и возраст угленосных отложений Чернышевского бурого угольного месторождения (Южное Приморье) // Тихоокеанская геология. № 4. С. 101–105].
- Bolotnikova, T.N. 1989. Age and correlation of coal-bearing strata of Rettikhovskii and Chernyshevskii brown coal field of the Southern Primorye (according to pollen data). In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 101–109, Dal'nevostochnoe otdelenie Akademii nauk SSSR, Vladivostok (in Russian). [Болотникова Т.Н. 1989. Возраст и корреляция угленосных толщ Реттиховского и Чернышевского бурого угольных месторождений Южного Приморья (по палинологическим данным) // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 101–109].
- Bolotnikova, T.N. 1993. Palynological characteristics of coal-bearing sediments of the Eastern section of the Pavlovskii brown coal field. *Tikhoookeanskaya geologiya* 6: 102–110. (in Russian). [Болотникова Т.Н. 1993. Палинологическая характеристика угленосных отложений Восточного участка Павловского месторождения // Тихоокеанская геология. № 6. С. 102–110].
- Bolotnikova, T.N. 1994. Palynostratigraphy of Cenozoic sediments of the Pavlovskii brown coal field. *Tikhoookeanskaya geologiya* 1: 71–81 (in Russian). [Болотникова Т.Н. 1994. Палиностратиграфия кайнозойских отложений Павловского бурого угольного месторождения // Тихоокеанская геология. № 1. С. 71–81].
- Bondarenko, O.V., N.I. Blokhina, A.A. Bruch, A.J. Henrot & T. Utescher 2017. Quantification of Calabrian vegetation in Southern Primorye (Far East of Russia) using multiple proxies. *Palaeogeography, Palaeoclimatology, Palaeoecology* 467: 253–264.
- Bondarenko, O.V., N.I. Blokhina, V. Mosbrugger & T. Utescher (in print). Paleogene climate dynamics in the Primorye Region, Far East of Russia, based on a Co-existence Approach analysis of palaeobotanical data. *Palaeobiodiversity and Palaeoenvironments*.
- Borsuk, M.O. 1952. *Fossil flora of the Upper Cretaceous sediments of Primorye*. Gosgeoltekhizdat, Moskva. 52 pp. (in Russian). [Борсук М.О. 1952. Ископаемая флора верхнемеловых отложений Приморья. Москва: Госгеолтехиздат. 52 с.].
- Chatshin, A.A., V.K. Popov, V.P. Nechaev, I.Yu. Chekryzhov, M.G. Nechaeva & M.G. Blokhin 2013. Geochemical features of the Eocene volcanism of rift basins of the South-Western Primorye. In: *Continental rifting, related processes* (S.V. Rasskazov, A.V. Nikishin, & S.P. Primina, eds.), pp. 164–168, Institut zemnoi kory Sibirskogo otdeleniya Rossiiskoi akademii nauk, Irkutsk (in Russian). [Чащин А.А., Попов В.К., Нечаев В.П., Чекрыжов И.Ю., Нечаева М.Г., Блохин М.Г. 2013. Геохимические особенности эоценового вулканизма рифтовых впадин Юго-Западного Приморья // Континентальный рифтогенез, сопутствующие процессы / под ред. С.В. Рассказова, А.М. Никишина, С.П. Приминной. Иркутск: Институт земной коры СО РАН. С. 164–168].
- Chekryzhov, I.Yu., V.K. Popov, A.M. Panichev, V.V. Seredin & E.V. Smirnova 2010. New data on stratigraphy, volcanism, and zeolite mineralization of the Cenozoic Vanchinskii Basin, Primorsky Krai. *Tikhoookeanskaya geologiya* 29(4): 45–63. (in Russian). [Чекрыжов И.Ю., Попов В.К., Паничев А.М., Середин В.В., Смирнова Е.В. 2010. Новые данные по стратиграфии, вулканизму и цеолитовой минерализации кайнозойской Ванчинской впадины, Приморский край // Тихоокеанская геология. Т. 29. № 4. С. 45–63].
- Cohen, K.M., S.M. Finney, P.L. Gibbard & J.X. Fan 2013. The ICS International Chronostratigraphic Chart. *Episodes* 36:199–204.
- Collinson, M.E. 1992. Vegetational and floristic changes around the Eocene/Oligocene boundary in western and central Europe. In: *Eocene-Oligocene climatic and biotic evolution* (D.R. Prothero & W.A. Berggren, eds.), pp. 437–450, Princeton University Press, Princeton.
- Dupont-Nivet, G., W. Krijgsman, C. Langereis, H.A. Abels, S. Dai & X. Fang 2007. Tibetan plateau aridification linked to global cooling at the Eocene–Oligocene transition. *Nature* 445:635–638.
- Eldrett, J.S., D.R. Greenwood, I.C. Harding & M. Huber 2009. Increased seasonality through the Eocene to Oligocene transition in northern high latitudes. *Nature* 459: 969–973.
- Flerov, K.K., E.I. Belyaeva, N.M. Yanovskaya, A.A. Gureev, I.M. Novodvorskaya, V.S. Kornilova, N.S. Shevyreva, E.N. Kurochkin, V.V. Zherikhin, V.M. Chkhikvadze, G.G. Martinson, N.V. Tolstikova, A.L. Chepalyga & L.I. Fofyanova 1974. *Zoogeography of the Paleogene of Asia* (Trudy Paleontologicheskogo instituta Akademii nauk SSSR, Vol. 146). Nauka, Moskva, 301 pp. (in Russian). [Флёров К.К., Беляева Е.И., Яновская Н.М., Гуреев А.А., Новодворская И.М., Корнилова В.С., Шевырёва Н.С., Курочкин Е.Н., Жерихин В.В., Чхиквадзе В.М., Мартинсон Г.Г., Толстикова Н.В., Чепалыга А.Л., Фотьянова Л.И. 1974.

- Зоогеография палеогена Азии (Труды ПИН АН СССР. Вып. 146). Москва: Наука. 301 с.].
- Gerasimov, I.P. (ed.) 1969. *Natural conditions and natural resources of the USSR. South Far East*. Nauka, Moskva. 424 pp. (in Russian). [Герасимов И.П. (ред.) 1969. Природные условия и естественные ресурсы СССР. Южная часть Дальнего Востока. Москва: Наука. 424 с.].
- Greenwood, D.R. & S.L. Wing 1995. Eocene continental climates and latitudinal temperature gradients. *Geology* 23: 1044–1048.
- Gromova, N.S. 1980. Palinocomplex of the Nazimovskaya formation of the Southern Primorye. Paleoclimatic studies for stratigraphy. *Trudy Vsesoyuznogo geologicheskogo instituta. Nov. ser.* 305: 80–83. (in Russian). [Громова Н.С. 1980. Палинокомплекс назимовской свиты Южного Приморья. Палеоклиматические исследования для целей стратиграфии // Труды ВСЕГЕИ. Нов. сер. № 305. С. 80–83].
- Klimova, R.S. 1981. New species of *Acer* from the Miocene of Northeastern Primorye. *Paleontologicheskii Zhurnal* 1: 134–138 (in Russian). [Климова Р.С. 1981. Новый вид *Acer* из миоцена Северо-Восточного Приморья // Палеонтологический журнал. № 1. С. 134–138].
- Klimova, R.S. 1988. Some representatives of heat-loving plants from the Miocene of Primorsky Krai. *Paleontologicheskii Zhurnal* 1: 92–99. (in Russian). [Климова Р.С. 1988. Некоторые представители теплолюбивых растений из миоцена Приморского края // Палеонтологический журнал. № 1. С. 92–99].
- Klimova, R.S., A.F. Kramchanin & T.I. Demidova 1977. New data on the stratigraphy of the Rettihovskii coal-bearing mine. In: *Stratigraphy of the Cenozoic sediments of the Far East* (Yu.A. Kosygin, ed.), pp. 66–75, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Климова Р.С., Крамчанин А.Ф., Демидова Т.И. 1977. Новые данные по стратиграфии Реттиховского угленосного разреза // Стратиграфия кайнозойских отложений Дальнего Востока / под ред. Ю.А. Косыгина. Владивосток: ДВНЦ АН СССР. С. 66–75].
- Klimova, R.S. & E.I. Tsar'ko 1989. On the discovery of Paleogene plants and diatoms in the Salibezskaya series of the Svetlovodnaya River basin (Northeastern Primorye). In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 57–63, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Климова Р.С., Царько Е.И. 1989. О находке палеогеновых растений и диатомей в салибезской толще бассейна реки Светловодная (Северо-Восточное Приморье) // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 57–63].
- Koshman, M.M. 1964. Tertiary flora of the Bikinskii brown coal field. *Botanicheskii Zhurnal* 49(2):265–271 (in Russian). [Кошман М.М. 1964. Третичная флора Бикинского бурогоугольного месторождения // Ботанический журнал. Т. 49. № 2. С. 265–271].
- Kovar-Eder, J. & Z. Kvaček 2003. Towards vegetation mapping based on the fossil plant record. *Acta Universitatis Carolinae, Geologica* 46(4):7–13.
- Kovar-Eder, J. & Z. Kvaček 2007. The integrated plant record (IPR) to reconstruct Neogene vegetation: the IPR-vegetation analysis. *Acta Palaeobotanica* 47(2):391–418.
- Kovar-Eder, J., H. Jechorek, Z. Kvaček & V. Parashiv 2008. The Integrated Plant Record: an essential tool for reconstructing Neogene zonal vegetation in Europe. *Palaios* 23: 97–111.
- Kovar-Eder, J. & V. Teodoridis 2018. The middle Miocene central European plant record revisited widespread sub-humid sclerophyllous forests indicated. *Fossil Imprint* 74 (1–2):115–134.
- Krassilov, V.A. 1972. *Paleoecology of terrestrial plants (basic principles and methods)*. Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok. 210 pp. (in Russian). [Красилов В.А. 1972. Палеоэкология наземных растений (основные принципы и методы). Владивосток: ДВНЦ АН СССР. 210 с.].
- Krassilov, V.A., 1989a. Cenozoic of the Far East: geodynamic scheme, reference section sections, florogenesis. In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 23–33, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Красилов В.А. 1989а. Кайнозой Дальнего Востока: геодинамическая схема, опорные разрезы, флорогенез // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 23–33].
- Krassilov, V.A., 1989b. Change of flora on the border of Cretaceous and Paleogene in the Kavalerskii District, Primorye. In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 34–37, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Красилов В.А. 1989б. Смена флоры на границе мела и палеогена в Кавалеровском районе, Приморье // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 34–37].
- Kundyshev, A.S. & T.I. Petrenko 1987. About the age of the Bolotninskaya fossil flora of the Southern Primorye. In: *Palinology of the East of the USSR* (V.S. Markevich, ed.), pp. 53–59, Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Кундышев А.С., Петренко Т.И. 1987. О возрасте болотнинской ископаемой флоры Южного Приморья // Палинология Востока СССР / под ред. В.С. Маркевич. Владивосток: ДВНЦ АН СССР. 53–59].
- Kundyshev, A.S. & N.B. Verkhovskaya 1989. About the age of coal-bearing sediments of the Nizhnebikinskii Basin. In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 121–127, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Кундышев А.С., Верховская Н.Б. 1989. О возрасте угленосных отложений Нижнебикинской впадины // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 121–127].
- Kurentsova, G.E. 1968. *Vegetation of Primorsky Krai*. Dal'nevostochnoe knizhnoe izdatel'stvo, Vladivostok. 192 pp. (in Russian). [Куренцова Г.Э. 1968. Растительность Приморского края. Владивосток: Дальневосточное книжное издательство. 192 с.].
- Mikhailov, V.A., Yu.M. Feoktistov & R.S. Klimova 1989. New data on the Cenozoic phytogeography of the eastern part of the Zerkal'enskii Basin. In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 38–49, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Михайлов В.А., Феоктистов Ю.М., Климова Р.С. 1989. Новые данные по фитогеографии кайнозой восточной части Зеркальненской депрессии // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 38–49].
- Oleinikov, A.V. & R.S. Klimova 1977. New data on stratigraphy of Neogene volcanogenic sediments of the

- Samarga River basin. In: Stratigraphy of the Cenozoic sediments of the Far East (Varnavskii V.G., ed.), pp. 76–80. Dal'nevostochnyi nauchnyi tsentr Akademii nauk SSSR, Vladivostok (in Russian). [Олейников А.В., Климова Р.С. 1977. Новые данные по стратиграфии неогеновых вулканогенных отложений бассейна реки Самарга // Стратиграфия кайнозойских отложений Дальнего Востока / под ред. В.Г. Варнавского. Владивосток: ДВНЦ АН СССР. С. 76–80].
- Pagani, M., J.C. Zachos, K.H. Freeman, B. Tipler & S. Bohlen 2005. Marked decline in atmospheric carbon dioxide concentrations during the Paleogene. *Science* 309: 600–603.
- Pavlyutkin, B.I. 2007a. *The Eocene us'davydovka flora of the south Primorye*. Dal'nauka, Vladivostok. 148 pp. (in Russian). [Павлюткин Б.И. 2007а. Эоценовая усть-давыдовская флора юга Приморья. Владивосток: Дальнаука. 148 с.].
- Pavlyutkin, B.I. 2007b. Representatives of the Ericaceae Superorder from the Oligocene Voznovskaya flora of the Primorskii Region. *Botanicheskii Zhurnal* 92(2):293–307 (in Russian). [Павлюткин Б.И. 2007б. Представители надпорядка Ericaceae из олигоценовой возновской флоры Приморского края // Ботанический журнал. Т. 92. № 2. С. 293–307].
- Pavlyutkin, B.I. 2011. The new species of thermophilic plants from the Early Oligocene of Kraskino (Primorsky Region) as evidence of its subtropic type. *Paleontologicheskii Zhurnal* 6:100–106. (in Russian). [Павлюткин Б.И. 2011. Новые виды термофильных растений в раннеолигоценовой флоре Краскино (Приморский край) как подтверждение ее субтропического облика // Палеонтологический журнал. № 6. С. 100–106].
- Pavlyutkin, B.I. & T.I. Petrenko 1993 New materials on the stratigraphy of Tertiary deposits of the Rechnoi Peninsula (Southern Primorye). *Tikhookeanskaya geologiya* 5: 42–50 (in Russian). [Павлюткин Б.И., Петренко Т.И. 1993. Новые материалы по стратиграфии третичных отложений полуострова Речной (Южное Приморье) // Тихоокеанская геология. № 5. С. 42–50].
- Pavlyutkin, B.I. & T.I. Petrenko 1994. To stratigraphy of the Tertiary coal-bearing sediments of the southeastern margin of the Khanka massif. *Tikhookeanskaya geologiya* 2: 18–29 (in Russian). [Павлюткин Б.И., Петренко Т.И. 1994. К стратиграфии третичных угленосных отложений юго-восточной окраины Ханкайского массива // Тихоокеанская геология. № 2. С. 18–29].
- Pavlyutkin, B.I. & T.I. Petrenko 1997. Problems of stratigraphy of Tertiary formations of Posyet Peninsula and the adjacent territory (South-West Primorye). *Tikhookeanskaya geologiya* 16(1): 89–98 (in Russian). [Павлюткин Б.И., Петренко Т.И. 1997. Проблемы стратиграфии третичных образований полуострова Посыет и прилегающей территории (Юго-Западное Приморье) // Тихоокеанская геология. Т. 16, № 1. С. 89–98].
- Pavlyutkin, B.I. & T.I. Petrenko 2010. *Stratigraphy of Paleogene – Neogene sediments in Primorye*. Dal'nauka, Vladivostok. 164 pp. (in Russian). [Павлюткин Б.И., Петренко Т.И. 2010. Стратиграфия палеоген – неогеновых отложений Приморья. Владивосток: Дальнаука. 164 с.].
- Pavlyutkin, B.I., S.I. Nevolina, T.I. Petrenko & T.K. Kutubzade 2006. On the age of the Paleogene Nazimovskaya and Khasanskaya Formations of the South-Western Primorye. *Stratigraphy and Geological Correlation* 14(3): 116–129 (in Russian). [Павлюткин Б.И., Неволина С.И., Петренко Т.И., Кутуб-Задэ Т.К. 2006. О возрасте палеогеновых назимовской и хасанской свит Юго-Западного Приморья // Стратиграфия и геологическая корреляция. Т. 14, № 3. С. 116–129].
- Pavlyutkin, B.I., I.Yu. Chekryzhov & T.I. Petrenko 2012. *Geology and flora of lower Miocene in the south Primorye*. Dal'nauka, Vladivostok. 194 pp. (in Russian). [Павлюткин Б.И., Чекрыжов И.Ю., Петренко Т.И. 2012. Геология и флора нижнего миоцена юга Приморья. Владивосток: Дальнаука. 194 с.].
- Pavlyutkin, B.I., I.Yu. Chekryzhov & T.I. Petrenko 2014. *Geology and floras of lower Oligocene in the Primorye*. Dal'nauka, Vladivostok. 236 pp. (in Russian). [Павлюткин Б.И., Чекрыжов И.Ю., Петренко Т.И. 2014. Геология и флора нижнего олигоцена Приморья. Владивосток: Дальнаука. 236 с.].
- Pearson, P.N., G.L. Foster & B.S. Wade 2009. Atmospheric carbon dioxide through the Eocene–Oligocene climate transition. *Nature* 461: 1110–1113.
- Popov, V.K., S.O. Maksimov, A.A. Vrzhosek & V.M. Chubarov 2007. Basaltoids and carbonatite tuffs of Amginskii volcano (South-Western Primorye): geology and genesis. *Tikhookeanskaya geologiya* 26(4):75–97 (in Russian). [Попов В.К., Максимов С.О., Вржосек А.А., Чубаров В.М. 2007. Базальтоиды и карбонатитовые туфы Амгинского вулкана (Юго-Западное Приморье): геология и генезис // Тихоокеанская геология. Т. 26, № 4. С. 75–97].
- Popov, V.K., S.V. Rasskazov, Yu.I. Chekryzhov et al. 2005. Potassium-argon dating and geochemical characteristics of trachybasalts and trachyandesites of Primorye. In: *Alkaline Earth Magmatism* (L.N. Kogarko, ed.), pp. 133–135, Institut Geokhimii i analiticheskoi khimii imeni V.I. Vernadskogo Rossiiskoi akademii nauk, Moscow (in Russian). [Попов В.К., Рассказов Ю.И., Чекрыжов Ю.И., и др. 2005. Калий-аргоновые датировки и геохимические характеристики трахибазальтов и трахиандезитов Приморья // Щелочной магматизм Земаи / под ред. Л.Н. Когарко. Москва: ГЕОХИ РАН. С. 133–135].
- Popova, S., T. Utescher, D.V. Gromyko, V. Mosbrugger, E. Herzog & L. François 2013. Vegetation change in Siberia and the northeast of Russia during the Cenozoic Cooling: a study based on diversity of plant functional types. *Palaio* 28: 418–432.
- Quan, C., C. Liu & T. Utescher 2012a. Eocene monsoon prevalence over China: A palaeobotanical perspective. *Palaogeography, Palaeoclimatology, Palaeoecology* 365–366: 302–311.
- Quan, C., C. Liu & T. Utescher 2012b. Paleogene temperature gradient, seasonal variation and climate evolution of northeast China. *Palaogeography, Palaeoclimatology, Palaeoecology* 313–314: 150–161.
- Rybalko, V.I., V.N. Ovechkin & R.S. Klimova 1980. Cenozoic basaltoids of the Amginskaya series (North-Eastern Primorye). *Sovetskaya geologia* 12:59–71 (in Russian). [Рыбалко В.И., Овечкин В.Н., Климова Р.С. 1980. Кайнозойские базальтоиды амгинской серии (Северо-Восточное Приморье) // Советская геология. № 12. С. 59–71].
- Sedova, M.A. 1957. Oligocene and Miocene floras and vegetation of the Primorskii Region according to the data of palynological analysis. In: *Issues of palaeobiogeography and biostratigraphy* (D.L. Stepanov, ed.), pp. 203–210, Gosgeoltekhizdat, Moscow (in Russian). [Седова М.А. 1957. Олигоценовые и миоценовые флоры и растительность Приморского края по данным палинологического анализа // Вопросы палеобио-

- географии и биостратиграфии / под ред. Д.А. Степанов. Москва: Госгеолтехиздат. С. 203–210].
- Tashchi, S.M., A.G. Ablaev & N.G. Mel'nikov 1996. *Cenozoic basin of the Western Primorye and adjacent territories of China and Korea*. Dal'nauka, Vladivostok. 168 pp. (in Russian). [Тащи С.М., Аблаев А.Г., Мельников Н.Г. 1996. Кайнозойский бассейн Западного Приморья и сопредельных территорий Китая и Кореи. Владивосток: Дальнаука. 168 с.].
- Teodoridis, V., J. Kovar-Eder, P. Marek, Z. Kvaček & P. Mazouch 2011. The integrated plant record vegetation analysis: internet platform and online application. *Acta Musei Nationalis Pragae, Ser. B* 67(3–4):159–165.
- Utescher, T., O.V. Bondarenko & V. Mosbrugger 2015. The Cenozoic Cooling – continental signals from the Atlantic and Pacific side of Eurasia. *Earth and Planetary Science Letters* 415:121–133.
- Utescher, T., A.A. Bruch, B. Erdei, L. François, D. Ivanov, F.M.B. Jacques, A.K. Kern, C. Liu, V. Mosbrugger & R.A. Spicer 2014. The coexistence approach – theoretical background and practical considerations of using plant fossils for climate quantification. *Palaeogeography, Palaeoclimatology, Palaeoecology* 410:58–73.
- Utescher, T., A.A. Bruch, A. Micheels, V. Mosbrugger & S. Popova 2011. Cenozoic climate gradients in Eurasia – a palaeo-perspective on future climate change? *Palaeogeography, Palaeoclimatology, Palaeoecology* 304: 351–358.
- Utescher, T., V. Mosbrugger, D. Ivanov & D.L. Dilcher 2009. Present-day climatic equivalents of European Cenozoic climates. *Earth and Planetary Science Letters* 284 (3–4):544–552.
- Varnavskii, V.G., A.K. Sedykh & V.I. Rybalko 1988. *Paleogene and Neogene Priamurye and Primorye*. Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok. 184 pp. (in Russian). [Варнавский В.Г., Седых А.К., Рыбалко В.И. 1988. Палеоген и неоген Приамурья и Приморья. Владивосток: ДВО АН СССР. 184 с.].
- Verkhovskaya, N.B. & A.S. Kundyshev 1989. Physiognomic features of spore-pollen spectra and their use in stratigraphy. In: *Cenozoic of the Far East* (V.A. Krassilov & R.S. Klimova, eds.), pp. 128–134, Dal'nevostochnoe otделение Akademii nauk SSSR, Vladivostok (in Russian). [Верховская Н.Б., Кундышев А.С. 1989. Физиономические особенности спорово-пыльцевых спектров и их использование в стратиграфии // Кайнозой Дальнего Востока / под ред. В.А. Красилова, Р.С. Климовой. Владивосток: ДВО АН СССР. С. 128–134].
- Xiao, G.Q., H.A. Abels, Z. Yao, G. Dupont-Nivet & F.J. Hilgen 2010. Asian aridification linked to the first step of the Eocene–Oligocene climate Transition (EOT) in obliquity-dominated terrestrial records (Xining Basin, China). *Climate of the Past* 6: 501–513.
- Yanovskaya, N.M. 1954. A new genus *Embolotheriidae* from the Paleogene of Mongolia. Tertiary mammals. *Trudy Paleontologicheskogo instituta Akademii nauk SSSR* 55(3): 5–43 (in Russian). [Яновская Н.М. 1954. Новый род *Embolotheriidae* из палеогена Монголии. Третичные млекопитающие // Труды ПИН АН СССР. Т. 55, № 3. С. 5–43].
- Zachos, J.C., G.R. Dickens & R.E. Zeebe 2008. An Early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature* 451:279–283.